

New Born Care: Our Perspective

BENJAMIN M SAGAYARAJ, NIDHI SHARMA, LAL D. V.

ABSTRACT

A low birth weight baby is a clinical and diagnostic challenge. The neonatologists are faced with numerous neonatal intensive care unit protocols. This study was designed to review the literature of past twenty five years regarding management of neonatal jaundice, sepsis, anaemia, hypoglycaemia, jaundice and hypoxic encephalopathy in a low birth weight newborn. The low birth weight newborn baby should be intubated electively if signs of respiratory distress appear. There should be an early Doppler of cerebral arteries to predict the ischaemic changes in neonatal brain. Probiotic therapy with *Bifidobacterium bifidus* and *Streptococcus*

thermophilus protects against necrotizing enterocolitis and results in increased weight gain. Newborn intravenous lines should not be flushed with normal saline ampoules containing benzyl alcohol as preservative, as this increases the fluidity of neonatal blood brain barrier and predisposes to neonatal jaundice. Erythropoietin subcutaneous injections are most rewarding in low birth weight babies with neonatal anaemia. There is also increase in weight. Kangaroo care is useful in management of neonatal hypothermia and is also an immunological boost as the baby gets colonised with favorable microorganisms of maternal skin.

Keywords: Anaemia, Hypothermia, Intensive care, Jaundice, Low birth weight, Neonatal

INTRODUCTION

Low birth weight may be the outcome of either preterm delivery (before 37 weeks of gestation) or retarded foetal (intrauterine) growth [1]. The problems of the intrauterine growth retarded babies are due to uteroplacental insufficiency and inadequate substrata transfer leading to birth asphyxia, hypothermia, meconium aspiration, polycythemia, hypoglycemia, hypocalcaemia, thrombocytopenia. On the other hand, the preterm complications are caused by anatomic and physiological immaturity. There is respiratory distress due to delayed alveolar clearance of alveolar fluid and surfactant deficiency. Postnatal circulatory adaptation is delayed due to pulmonary hypertension, systemic hypotension and delayed closure of the three foetal shunts (ductus venosus, ductus arteriosus and foramen ovale). Liver immaturity and reduced substrate explain the high prevalence of jaundice and deficiency of coagulation factors. Immature vascular development in retina and central nervous system predisposes to retinopathy and intraventricular haemorrhage. Immature skin and mucosa barrier and immature cellular and humoral immunity predisposes to neonatal sepsis and nosocomial infections. Gastrointestinal mucosa is immunologically immature. Besides this there is inappropriate bacterial gut colonisation, this results in necrotizing enterocolitis.

A growth restricted preterm baby is (clinically and diagnostically a challenge) because the dual problems of retardation and prematurity are superimposed.

The proportion of low birth weight has increased in the past twenty years. This increase is attributed to changes in the frequency of multiple births, increase in obstetrical interventions, and improved ascertainment of early preterm births and increased use of ultrasound for estimation of gestational age. Unfortunately the documentation of low birth weight babies is not proper [2].

There has been considerable advances in neonatal care in past 25 years. However, low birth weight babies have an altogether altered developmental, metabolic, and nutritional status. Routine NICU protocols have to be modified for these babies with special needs [3-8].

LOW BIRTH WEIGHT PROBLEMS

The risk to a low birth weight newborn baby can be classified as:

- 1) **Early risk:** In the delivery room and neonatal life. The nursery neurobiological events of a low birth weight baby are incidence of neonatal septicemia, low blood pH, seizures, intraventricular leukomalacia and hypoglycaemia. These factors correlate well with deficits in Bayley scale of infant development, mental developmental index and psychomotor development index [9]. Relative importance of individual factors can be assessed as neurobiological risk score. This score correlates well with mental, motor, neurological outcomes [10].
- 2) **Late risk:** Long term developmental outcomes. The increased survival of low birth weight babies



[Table/Fig-1]: A preterm baby in an ultraviolet light double surface phototherapy unit

has resulted in an increased incidence of cerebral palsy. There are two postulated hypotheses [11]. One theory is that as more low birth weight babies survive, they suffer the complications of extreme prematurity like periventricular leukomalacia, intraventricular haemorrhage, respiratory distress and sepsis resulting in postnatal brain damage in an otherwise normal infant. Another hypothesis is that cerebral palsy and preterm birth have similar pathophysiology in antenatal period and that these babies who were compromised well before birth are now surviving. Environmental enrichment programs are most effective in moderately low birth weight child who comes from a lower socio economic status [11].

CARE OF THE LOW BIRTH WEIGHT BABIES

Appropriate resuscitation/ respiratory care

Perinatal asphyxia is recognised as a major cause of neonatal morbidity and mortality in the developing countries. The effects are more pronounced among low birth weight babies. Delivery room policies have changed extremely. Until 1994, the low birth weight infants were intubated immediately after delivery, when presenting the slightest signs of respiratory distress or asphyxia. Later the guidelines were to do the continuous (15-20 seconds) pressure control (20-25cm H₂O) inflation of the lungs using a nasal pharyngeal tube, followed by continuous positive airway pressure (4 to 6 cm H₂O) applied to all extremely low birth weight immediately after delivery to establish a functional residual capacity and perhaps to avoid elective intubation and mechanical ventilation [12]. Literature search suggested no difference in the initial ventilator settings, ventilator days. The mortality and morbidity rates were found similar in extremely low birth weight babies with primary endotracheal intubation and mechanical ventilation in the delivery room as compared to infants with secondary endotracheal intubation and mechanical ventilation attributed to respiratory distress syndrome later in the intensive care unit.

Extremely low birth weight babies which did not require endotracheal intubation and mechanical ventilation had no increased incidence of bronchopulmonary dysplasia or periventricular leukomalacia and intraventricular hemorrhage.

It was inferred that individualised intubation strategy is safe as compared to routine immediate intubation of all extremely low birth weight babies.

Neonatal hypoglycemia and Feeding

In low birth weight babies there is reduced reserve. Besides, increased utilisation during birth, hypoxia results in neonatal hypoglycemia. This is defined as a serum glucose level <30mg/100mg in the first few hours of birth. The highest incidence of hypoglycemia (67%) occurred in preterm low birth weight. It was 25% in term low birth weight babies and 18% in post term low birth weight babies [13].

Early initiation and frequent (2 to 3 hourly) feeding with breast milk is the most cost effective strategy in the prevention of hypoglycaemia. Breast milk promotes ketogenesis and has a lower insulinogenic effect [14, 15]. Alternative feeds can be utilised in maternal HIV infection. In preterm neonates fortification of breast milk is essential. Micronutrients supplementation with vitamins and minerals promote better assimilation and weight gain. Total parenteral nutrition can be continued till 100mL/kg/day, and is supplied by the enteral route. The amount of feeding is increased slowly and stopped if any signs of feeding intolerance (vomiting, abdominal distension and increased gastric residual volume) appear.

Prevention of Necrotising Enterocolitis

At birth the basic body tries to develop an intricate symbiotic equilibrium between bacterial environment and its own immune system-an equilibrium that results in the preferential colonization of the gastrointestinal tract by a variety of "favorable" gram positive microorganisms most notably *Lactobaccillus* and *Bifidobacterium*.

In contrast, the low birth weight new born's intestine tends to be colonized by coliforms, enterococci and bacteroids. *Bifidobactium* and *Lactobaccillus* are found in the stools of <5% extremely low birth infants within the first month of life [16].

Probiotics strengthen the intestinal mucosal barrier which impedes the translocation of pathogenic bacilli in IL-10 knocked out mice. It was postulated that probiotics decrease cytokine production both systemically and at mucosal surface [17].

The pathogenesis of necrotising enterocolitis is multi factorial. Prematurity, formula feeding, intestinal ischaemia and bacterial colonisation activate an inflammatory cascade accumulating in bowel necrosis [18]. We speculated to create fresh suspension (0.5

teaspoon) of probiotic powder (ABC Dophilus) diluted in 3ml of mother's milk daily along with regular feeds. This provides 1.05×10^9 colony forming units/day, consisting of 0.35×10^9 CFU *Bifidobacterium infantis*, 0.35×10^9 CFU *Streptococcus thermophilus* and 0.35×10^9 of *Bifidobacterium bifidus*. We added *S. thermophilus* as studies suggested the prevention of secondary rotavirus diarrhea when *S. thermophilus* was added [19]. We decided not to include Lactobacillus, as cases of lactobacillus bacteremia have been described in immune compromised babies receiving high doses of lactobacillus [20].

Prevention of Hypothermia

Conductive and evaporative heat loss can be prevented by warming of delivery rooms, immediate drying and wrapping after birth, frequent feeding, delayed bathing and kangaroo mother care. Kangaroo mother care provides additional benefits as lowered incidence of nosocomial infections, as the babies skin gets colonised with friendly maternal skin microbes. There are other advantages as well. There is less incidence of lower respiratory tract disease, promotion of exclusive breast feeding and better weight gain [21]. Incubators, semi permeable plastic sheets and baby closure are other useful methods to prevent hypothermia.

Infection control

Neonatal infection contributes to 8-80% of all neonatal deaths [21]. Approaches that have been used to prevent infection among low birth weight babies include strict policy of clean hands and avoiding overcrowding. Colonising babies with maternal organisms via kangaroo mother care improves immunological maturation. This also leads to a favorable bacterial colonisation [22]. Keeping the nursery equipment clean and on regular bacterial colony surveillance prevents nosocomial infections. Preparing fresh feeds and hygienic dedicated nursing staff prevents incidence of sepsis. Patients education and visitors policy are also carried out.

Neonatal Anaemia

Anaemia of prematurity is usually caused by inadequate erythropoietin production. In addition diagnostic tests cause blood loss.

We encourage delayed cord clamping, after the umbilical cord pulsations have stopped, in low birth weight babies. The corrected reticulocyte count is calculated by normalizing to a haematocrit of 45 percent by multiplying the measured reticount with actual haematocrit and dividing by 45.

Oral iron supplementation (2mg/kg) is started on day 14 in all infants. If the serum ferritin fall below 100ng/mL or clinical signs of iron deficiency appear, the dose of iron is increased and 250 IU of epoetin beta per kilogram, are given three times a week, subcutaneously after D3 of life till D42. Infants who were on ventilator or who were

less than two weeks old with signs of anaemia were transfused if their haematocrit fall below 40% or their haemoglobin concentration fall below 14gm per deciliter. If blood samples of total 9 mL/kg have been obtained since previous transfusion, a repeat transfusion was considered. In spontaneously breathing babies older than two weeks, if fraction of inspired oxygen was less than 0.40, and if they had clinical signs of anaemia or their haematocrit was less than 32% and hemoglobin less than 11 gm/dL transfusion was given. The heart rate, respiratory rate, blood pressure, blood gas values and renal and liver functions are monitored. In our NICU, blood sampling is done on day 3, day 12, day 24, day 40. The low birth weight babies, whose birth weight is more than 1200 gms and whose baseline haematocrit is more than 48% respond maximum to epoetin beta [23].

Intraventricular Haemorrhage and Retinopathy.

The cranial ultrasound scans are done on day 3 and day 12 to detect intraventricular haemorrhage if hypoxic ischaemic injury in extremely low birth weight neonates is suspected. The regional cerebral blood flow study with power and pulsed Doppler is recommended, to detect impaired cerebrovascular autoregulation, Measuring the resistivity index in lenticulostriate arteries, in first week of life, can identify preterm neonates who are at risk of developing periventricular leukomalacia or germinal matrix haemorrhage or both [24].

Ophthalmoscopy is done twice weekly, till 4 weeks after the estimated date of delivery, to detect retinopathy of prematurity at the earliest.

Hyperbilirubinemia

The hyperbilirubinemia in preterm infants is because of red cell, hepatic and gastrointestinal immaturity. After delivery, the postnatal hepatic bilirubin uptake and conjugation is slow. In preterm babies the delayed feeding limits intestinal flow and colonisation [25]. This results in increased bilirubin in enterohepatic circulation.

There should be no flushing of intravenous catheter with bacteriostatic saline ampules containing benzyl alcohol. Benzyl alcohol increases membrane fluidity and may facilitate the passage of bilirubin into the brain Aggressive phototherapy and exchange transfusions if total serum bilirubin is 20mg/dL [26] is recommended. It is unlikely that hyperbilirubinemia has a causal relationship with periventricular leukomalacia. Periventricular leukomalacia is primarily an ischaemic lesion, probably caused by hypoperfusion. Bilirubin is primarily toxic to neurons and not to glial elements which predominate in the periventricular white matter [25].

CONCLUSION

Neonatologists are in a clinical quandary with respect to NICU protocols. This study was designed to compare various articles on aggressive and conservative

management strategies for low birth weight babies. By articulating the various research publications in terms of peculiar metabolic and developmental requirements of low birth weight babies, an approach has been developed. Standardisation of clinical methods and educating the paramedical staff is important for the success of any medical intervention. Training of mid-level health personnel on appropriate care of the low birth weight and providing management protocols/algorithms is of great importance. Record keeping for care audit and ongoing research is important.

REFERENCES

- [1] United Nations Children Fund and World Health Organization. Low birth weight: country, regional and global estimates. *UNICEF*. 2004;92-806-3832-37.
- [2] Victoria CG, Wagstaff A, Schellenberg JA, Gwatkin D, Claeson M and Habicht JP. Applying an equality lens to child health and mortality: more of same is not enough. *The Lancet*. 2003; 362:233-41.
- [3] Fanaroff AA, Stoll BJ, Wright LL, Carlo WA, Ehrenkranz RA, Stark AR, et al. NICHD Neonatal research network. Trends in neonatal morbidity and mortality for very low birth weight infants. *Am J Obstet Gynecol*. 2007; 196(2):147. e1-8.
- [4] Horbar JD, Carpenter JH, Badger GJ, Kenny M J, Roger F Soll, Kate A, Morrow and Jeffrey S. Buzas. Mortality and Neonatal Morbidity among infants 501-1500gms from 2000-2009. *Pediatrics*. 2012; 129:1019-26.
- [5] Costeloe KL, Hennessy EM, Haider S, Stacey F, Neil Marlow, Elizabeth S Draper . Short term outcomes after extreme preterm birth in England: comparison of two birth weight cohorts in 1995 and 2006(the EPIcure studies) *BMJ*. 2012;345: e7976.
- [6] Luregn JS, Adams M, Proietti E, Aebischer M, Grunt S, Barradori-Tolsa C, et al. Outcome at two years of age in a swiss national cohort of extremely preterm infants between born between 2000 and 2008. *BMC Pediatrics*. 2012; 12:198.
- [7] Ruegger C, Hegglin M, Adams M, Hans Ulrich Bucher and Swiss neonatal network. Population based trends in mortality, morbidity and treatment for very preterm and very low birth weight infants over 12 years. *BMC Pediatrics*. 2012;12:17.
- [8] Shah PS , Ye XY, Synnes A, Rouvinez-Bouali N, Yee W, Lee SK and Canadian Neonatal Network. Prediction of survival without morbidity for infants born at under 33 weeks of gestation age: a user friendly graphical tool. *Arch Dis Child Fetal Neonatal Ed*. 2012; 97(2):F110-15.
- [9] Brazy JE, Eckerman CO, Oehler JM, Goldstein RI, O'Rand AN. Nursery neurobiologic risk score: Important factors in predicting outcome in very low birth weight infants. *The Journal of Pediatrics*. 1991; 118(5):783-92.
- [10] Maureen Hack, Nancy K. Klein, H. Gerry Taylor. Long term developmental outcomes of low birth weight infants. *JSTOR: The future of children*. 1995; 5(1):176-96.
- [11] Stanely FJ, Watson L. Trends in perinatal mortality and cerebral palsy in western Australia, 1967 to 1985. *BMJ*. 1992; 304:1658-63.
- [12] Lindner W, Vobbeck S, Hummler H, Pontandt F. Delivery room management of extremely low birth weight infants: Spontaneous breathing or Intubation? *Pediatrics*. 103 ;5:961-67.
- [13] Lubchenco LO, Bard H. Incidence of hypoglycemia in newborn infants classified by birth weight and gestational age. *Pediatrics*. 1971; 47(5):831-38.
- [14] Lucas A, Boyes S, Bloom S and Aynsley Green A . Metabolic and endocrine responses to a milk feed in six day old infants; differences between breast and cow's milk formula feeding. *Act a pediatrics*. 1981; 70: 195-200.
- [15] Howdon JM, World Plati MP, Aynsley Green A. Pattern of metabolic adaptation for preterm and term infants in the first neonatal week. *Arch Dis Child*. 1992; 67: 357-65.
- [16] Gewolb IH, Schwalbe RS, Taciak VL, Harrison TS, Panigrani P. Stool microflora in extremely low birth weight infants. *Arch Dis Child Fetal Neonatal*. 1999; 80:F167-73.
- [17] Mc Carthy J, O'Mahony L, O'Callaghan L, Sheil B, Vaughan EE, Fitzsimons N. Double blind placebo controlled trial of two probiotic strains in interleukin 10 knockout mice and mechanistic link with cytokine balance. *Gut*. 2003; 52:975-80.
- [18] Saavedra JM, Bauman NA, Oung I, Perman JA, Yolken RH. Feeding of Bifidobacterium bifidum and Streptococcus thermophilus to infants in hospital for prevention of diarrhoea and shedding of rotavirus. *Lancet*. 1994; 344:1046-49.
- [19] Nopchinda S, Varavithya W, Phuapradit P, Sangchai R, Suthutvoravut U, Chantraruksa V. Effect of Bifidobacterium Bb12 with or without Streptococcus thermophilus supplemented formula on nutritional status. *J Med Assoc Thai*. 2002,85 (suppl 4): S1225-31.
- [20] Cnota J, Sheety AK, Land MH, Rouster-Stevens K, Woods C, Cannon ML. Sepsis associated with probiotic therapy lactobacillus. *Pediatrics* 2005; 115:178-81.
- [21] Charpak N, Ruiz-Pelaez J, Figueroa C and Charpak Y. A randomized controlled trail of Kangaroo mother care: results of follow up at 1 year of corrected age. *Pediatrics*. 2001; 108:1072-79.
- [22] Thaver D. Burden of Neonatal infections in Developing countries: A Review of Evidence from Community Based Studies. *Pediatr Infect Dis J*. 2009; 28:53-59.
- [23] Maier RF, Obladen M, Linderkamp PSO, Duc G, Hieronimi G, Halliday HL, Hans T. Versmold, et al. *The New England Journal of Medicine*. 1994. 30;17:1173-78.
- [24] F G Blankenberg, N N Loh, A M Norbash, J A Craychee, D M Spielman, B L Person, C A Berg , D R Enzmann. Impaired cerebrovascular autoregulation after hypoxic-ischemic injury in extreme low birth weight neonates: detection with power and pulsed wave Doppler US. *RSNA Radiology*. 1997; 205(2).
- [25] J F Watchko, M J Maisels. Jaundice in low birth weight infants: pathobiology and outcome. *Arch Dis Child Fetal Neonatal Ed*. 2003;88:F455-F458.
- [26] Wennberg RP. Cellular basis of bilirubin toxicity. *N Y State J Med*. 1991; 91:493-502.

AUTHOR(S):

1. Dr. Benjamin M Sagayaraj
2. Dr. Nidhi Sharma
3. Dr. Lal D. V.

PARTICULARS OF CONTRIBUTORS:

1. Associate Professor, Department of Paediatrics, Saveetha Medical College, Saveetha Nagar, Thandalam, Chennai, India.
2. Associate Professor, Department of Obstetrics and Gynaecology, Saveetha Medical College, Saveetha Nagar, Thandalam, Chennai, India.
3. Senior Resident, Department of Paediatrics, Saveetha Medical College, Saveetha Nagar, Thandalam, Chennai, India.

NAME, ADDRESS, E-MAIL ID OF THE CORRESPONDING AUTHOR:

Dr. Benjamin M. Sagayaraj
Associate Professor
No 11/2 APPU Street , Second lane, Mylapore
Chennai 600004, India.
Phone: 9445560392
Email: getdrben@yahoo.com

FINANCIAL OR OTHER COMPETING INTERESTS:

None.

Date of Publishing: Oct 30, 2014